

LOCAL COHOMOLOGY BOUNDS AND THE WEAK IMPLIES STRONG CONJECTURE IN DIMENSION 4

IAN ABERBACH AND THOMAS POLSTRA

ABSTRACT. We verify the weak implies strong conjecture for 4-dimensional finitely generated algebras over a field of prime characteristic $p > 5$ which has infinite transcendence degree over \mathbb{F}_p .

1. INTRODUCTION

Let (R, \mathfrak{m}, k) be an excellent commutative Noetherian local ring of prime characteristic $p > 0$. Tight closure theory, introduced and developed by Hochster and Huneke in [HH89, HH90, HH91, HH93, HH94a, HH94b], is a subject central to prime characteristic commutative algebra, important to our understanding of characteristic 0 singularities, and a guiding force in the development of mixed characteristic singularities.

Let $N \subseteq M$ be R -modules and let N_M^* denote the tight closure of N inside of M , see [HH90] for a precise definition of tight closure. The ring R is *weakly F -regular* if $N_M^* = N$ for all finitely generated modules $N \subseteq M$ and R is *strongly F -regular* if $N_M^* = N$ for all R -modules $N \subseteq M$. The weak implies strong conjecture asserts that every weakly F -regular ring is strongly F -regular. The most notable cases that the weak implies strong conjecture has been proven for are for standard graded algebras over a field, [LS99], and all rings of Krull dimension at most 3, [Wil95].

Outside of the graded scenario, developments around the weak implies strong conjecture depend upon the behavior of the anticanonical algebra of R . An unpublished result of Singh asserts that the weak implies strong conjecture holds for the class of rings whose anticanonical algebra is Noetherian¹, see [CEMS18b, Corollary 5.9]. Williams' proof of the weak implies strong conjecture in dimension 3 implicitly utilizes that the anticanonical algebra of a 3-dimensional weakly F -regular ring is Noetherian on the punctured spectrum, an assertion that relies on the algebraic-geometric methods of [Lip69] and [Smi97].

Our methods show that every 4-dimensional weakly F -regular ring is strongly F -regular, provided its anticanonical algebra is Noetherian when localized at a non-maximal prime ideal. Therefore a further relationship between tight closure and prime characteristic birational geometry is desired to progress the weak implies strong conjecture. This is indeed the scenario in 4-dimensional rings. Recent developments in the prime characteristic minimal model program, [DW19], allow us to conclude that all divisorial blowups rings, including

Polstra was supported in part by NSF Grant DMS #101890 during the preparation of this article.

¹It is conjectured that any divisorial blowup ring, including the anticanonical algebra, of any strongly F -regular ring is Noetherian.

the anticanonical algebra, are Noetherian away from maximal ideals for large classes of 4-dimensional weakly F -regular rings.

Theorem A. *Let R be a 4-dimensional finitely generated algebra over a field of prime characteristic $p > 5$ which has infinite transcendence degree over \mathbb{F}_p . If R is weakly F -regular then R is strongly F -regular.*

The weak implies strong conjecture will be understood as a problem of understanding when an element of a non-injective direct limit system is mapped to 0 inside the direct limit. To this end, we introduce the notion of a *local cohomology bound* in Section 2. If M is an R -module and $\underline{x} = x_1, \dots, x_\ell$ a sequence of elements then we may identify the local cohomology module $H_{(\underline{x})}^i(M)$ as a direct limit of Koszul cohomologies

$$H_{(\underline{x})}^i(M) \cong \varinjlim_{t_1 \leq t_2} \left(H^i(x_1^{t_1}, \dots, x_\ell^{t_1}; M) \xrightarrow{\alpha_{t_1, t_2}^i} H^i(x_1^{t_2}, \dots, x_\ell^{t_2}; M) \right).$$

The i th local cohomology bound of M with respect to the sequence \underline{x} is bounded by an integer k if for every t , if $\eta \in H^i(x_1^t, \dots, x_\ell^t; M)$ represents the 0-element of $H_{(\underline{x})}^i(M)$ then $\alpha_{t, t+k}^i(\eta) = 0$. Understanding when a module has bounded local cohomology bounds with respect to a sequence of elements is an interesting, challenging, and worth-while venture.

2. LOCAL COHOMOLOGY BOUNDS

We do not present the basic theory of local cohomology bounds in full generality. We only present specific aspects needed in later sections. The interested reader should consult [AP21] for a thorough introduction to the theory of local cohomology bounds.

2.1. Definition of local cohomology bound. Suppose M is a module over a ring R and $\underline{y} = y_2, y_3, y_4$ ² a sequence of elements. Then for each integer $t \in \mathbb{N}$ we let $\underline{y}^t = y_2^t, y_3^t, y_4^t$ and for each pair of integers $t_1 \leq t_2$ let $\tilde{\alpha}_{M; \underline{y}; t_1; t_2}^\bullet$ denote the natural map of Koszul cocomplexes

$$K^\bullet(\underline{y}^{t_1}; M) \xrightarrow{\tilde{\alpha}_{M; \underline{y}; t_1; t_2}^\bullet} K^\bullet(\underline{y}^{t_2}; M).$$

²We begin the sequence at y_2 instead of y_1 for ease of referencing this material in Section 3 and Section 4.

More specifically, $\tilde{\alpha}_{M;\underline{y};t_1;t_2}^\bullet$ is the following map of Koszul cocomplexes:

(2.1)

$$\begin{array}{ccccccccc}
K^\bullet(\underline{y}^{t_1}; M) : & 0 & \longrightarrow & M & \xrightarrow{\begin{bmatrix} y_4^{t_1} \\ y_3^{t_1} \\ y_2^{t_1} \end{bmatrix}} & M^{\oplus 3} & \xrightarrow{\begin{bmatrix} y_3^{t_1} & -y_4^{t_1} & 0 \\ -y_2^{t_1} & 0 & y_4^{t_1} \\ 0 & y_2^{t_1} & -y_3^{t_1} \end{bmatrix}} & M^{\oplus 3} & \xrightarrow{\begin{bmatrix} y_2^{t_1} \\ y_1^{t_1} \\ y_4^{t_1} \end{bmatrix}} & M & \longrightarrow & 0 \\
\downarrow \tilde{\alpha}_{M;\underline{y};t_1;t_2}^\bullet & & & \downarrow \tilde{\alpha}_{M;\underline{y};t_1;t_2}^0 & & \downarrow \tilde{\alpha}_{M;\underline{y};t_1;t_2}^1 & & \downarrow \tilde{\alpha}_{M;\underline{y};t_1;t_2}^2 & & \downarrow \tilde{\alpha}_{M;\underline{y};t_1;t_2}^3 & & & \\
K^\bullet(\underline{y}^{t_2}; M) : & 0 & \longrightarrow & M & \xrightarrow{\begin{bmatrix} y_4^{t_2} \\ y_3^{t_2} \\ y_2^{t_2} \end{bmatrix}} & M^{\oplus 3} & \xrightarrow{\begin{bmatrix} y_3^{t_2} & -y_4^{t_2} & 0 \\ -y_2^{t_2} & 0 & y_4^{t_2} \\ 0 & y_2^{t_2} & -y_3^{t_2} \end{bmatrix}} & M^{\oplus 3} & \xrightarrow{\begin{bmatrix} y_2^{t_2} \\ y_1^{t_2} \\ y_4^{t_2} \end{bmatrix}} & M & \longrightarrow & 0
\end{array}$$

Where

- $\tilde{\alpha}_{M;\underline{y};t_1;t_2}^0 = \text{id}_M$;
- $\tilde{\alpha}_{M;\underline{y};t_1;t_2}^1 = \begin{bmatrix} y_4^{t_2-t_1} & 0 & 0 \\ 0 & y_3^{t_2-t_1} & 0 \\ 0 & 0 & y_2^{t_2-t_1} \end{bmatrix}$
- $\tilde{\alpha}_{M;\underline{y};t_1;t_2}^2 = \begin{bmatrix} (y_3y_4)^{t_2-t_1} & 0 & 0 \\ 0 & (y_2y_4)^{t_2-t_1} & 0 \\ 0 & 0 & (y_2y_3)^{t_2-t_1} \end{bmatrix}$
- $\tilde{\alpha}_{M;\underline{y};t_1;t_2}^3 = \cdot (y_2y_3y_4)^{t_2-t_1}$

We let $\alpha_{M;\underline{y};t_1;t_2}^j$ denote the induced map of Koszul cohomologies

$$H^j(\underline{y}^{t_1}; M) \xrightarrow{\alpha_{M;\underline{y};t_1;t_2}^j} H^j(\underline{y}^{t_2}; M).$$

In particular,

$$\varinjlim_{t_1 \leq t_2} \left(H^j(\underline{y}^{t_1}; M) \xrightarrow{\alpha_{M;\underline{y};t_1;t_2}^j} H^j(\underline{y}^{t_2}; M) \right) \cong H_{(\underline{y})}^j(M)$$

by [BH93, Theorem 3.5.6].

For each $0 \leq j \leq 3$ let $\alpha_{M;\underline{y};t;\infty}^j$ be the natural map

$$H^j(\underline{y}^t; M) \xrightarrow{\alpha_{M;\underline{y};t;\infty}^j} H_{(\underline{y})}^j(M).$$

An element $\eta \in H^j(\underline{y}^t; M)$ belongs to $\text{Ker}(\alpha_{M;\underline{y};t;\infty}^j)$ if and only if there exists some $k \geq 0$ so that $\eta \in \text{Ker}(\alpha_{M;\underline{y};t+k}^j)$. If $\eta \in \text{Ker}(\alpha_{M;\underline{y};t;\infty}^j)$ we let

$$\epsilon_{\underline{y},t}^j(\eta) = \min\{k \mid \eta \in \text{Ker}(\alpha_{M;\underline{y};t+k}^j)\}.$$

Definition 2.1. Let R be a ring, $\underline{y} = y_2, y_3, y_4$ a sequence of elements in R , and M an R -module. The j th local cohomology bound of M with respect to the sequence of elements \underline{y} is

$$\text{lcb}_j(\underline{y}; M) = \sup\{\epsilon_{\underline{y},t}^j(\eta) \mid \eta \in \text{Ker}(\alpha_{M;\underline{y};t;\infty}^j) \text{ for some } t\} \in \mathbb{N} \cup \{\infty\}.$$

Observe that if M is an R -module and $\underline{y} = y_2, y_3, y_4$ a sequence of elements, then $\text{lcb}_j(\underline{y}; M) \leq N < \infty$ simply implies that if $\eta \in H^j(\underline{y}^t; M)$ represents the 0-element in the direct limit

$$\varinjlim_{t_1 \leq t_2} \left(H^j(\underline{y}^{t_1}; M) \xrightarrow{\alpha_{M; \underline{y}; t_1; t_2}^j} H^j(\underline{y}^{t_2}; M) \right) \cong H_{(\underline{y})}^j(M)$$

then $\alpha_{M; \underline{y}; t; t+N}^j(\eta)$ is the 0-element of the Koszul cohomology group $H^j(\underline{y}^{t+N}; M)$. Therefore finite local cohomology bounds correspond to a uniform bound of annihilation of zero elements in a choice of direct limit system defining a local cohomology module.

2.2. Some basic properties of local cohomology bounds.

Lemma 2.2. *Let R be a commutative Noetherian ring, M an R -module, and $\underline{y} = y_2, y_3, y_4$ a sequence of elements, then $\text{lcb}_j(\underline{y}^t; M) \leq \text{lcb}_j(\underline{y}; M)$. Furthermore, $\text{lcb}_j(\underline{y}; M) \leq tm$ for some integers t, m if and only if $\text{lcb}_j(\underline{y}^t; M) \leq m$.*

Proof. One only has to observe that $\alpha_{M; \underline{y}^t; k; k+m}^j = \alpha_{M; \underline{y}; tk; tk+tm}^j$. \square

Proposition 2.3. *Let R be a commutative Noetherian ring and M, N modules over R . Suppose $\underline{y} = y_2, y_3, y_4$ is a sequence of elements so that $(y_2, y_3, y_4)M = 0$ and $(y_2, y_3)N = 0$. Then*

- (1) $\alpha_{M; \underline{y}; t; t+k}^j = 0$ for all $t, k \geq 1$, and $1 \leq j \leq 3$;
- (2) $\alpha_{N; \underline{y}; t; t+k}^j = 0$ for all $t, k \geq 1$, and $2 \leq j \leq 3$.

In particular, $\text{lcb}_j(\underline{y}; M) \leq 1$ for $1 \leq j \leq 3$ and $\text{lcb}_j(\underline{y}; N) \leq 1$ for $2 \leq j \leq 3$.

Proof. Recall that $\alpha_{M; \underline{y}; t; t+k}^j$ is the map of Koszul cohomologies induced from the map $\tilde{\alpha}_{M; \underline{y}; t; t+k}^\bullet$ on Koszul cocomplexes. One can consult the diagram of (2.1) to observe that $\tilde{\alpha}_{M; \underline{y}; t; t+k}^j$ is the 0-map for all $t, k \geq 1$, and $j \geq 1$. The second assertion follows by an identical argument. \square

Proposition 2.4. *Let (R, \mathfrak{m}, k) be a local ring and*

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$$

a short exact sequence of finitely generated R -modules. Let $\underline{y} = y_2, y_3, y_4$ a sequence of elements of R . If $(y_2, y_3)M_3 = 0$ then

$$\text{lcb}_3(y_2, y_3, y_4; M_1) \leq \text{lcb}_3(y_2, y_3, y_4; M_2) + 1.$$

Proof. Consider the following commutative diagram, whose middle row is exact:

$$\begin{array}{ccccc} & & H^3(\underline{y}^t; M_1) & \longrightarrow & H^3(\underline{y}^t; M_2) \\ & & \downarrow \alpha_{M_1; \underline{y}; t; t+k}^3 & & \downarrow \alpha_{M_2; \underline{y}; t; t+k}^3 \\ H^2(\underline{y}^{t+k}; M_3) & \longrightarrow & H^3(\underline{y}^{t+k}; M_1) & \longrightarrow & H^3(\underline{y}^{t+k}; M_2) \\ & & \downarrow \alpha_{M_3; \underline{y}; t+k; t+k+1}^2 & & \downarrow \alpha_{M_1; \underline{y}; t+k; t+k+1}^3 \\ H^2(\underline{y}^{t+k+1}; M_3) & \longrightarrow & H^3(\underline{y}^{t+k+1}; M_1) & & \end{array}$$

By Proposition 2.3 the map $\alpha_{M_1; \underline{y}; t+k; t+k+1}^2$ is the 0-map. A straightforward diagram chase of the above diagram, which follows an element $\eta \in \text{Ker}(\alpha_{M_2; \underline{y}; t; t+k'}^3)$ for some k' , shows that $\eta \in \text{Ker}(\alpha_{M_2; \underline{y}; t; t+k+1}^3)$ whenever $k \geq \text{lcb}_3(\underline{y}; M_3)$. In particular, $\text{lcb}_3(\underline{y}; M_2) \leq \text{lcb}_3(\underline{y}; M_3) + 1$. \square

Proposition 2.5. *Let R be a commutative Noetherian ring, $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ a short exact sequence of R -modules, and $\underline{y} = y_2, y_3, y_4$ a sequence of elements in R .*

- (1) *If \underline{y} is a regular sequence on M_2 then $\text{lcb}_j(\underline{y}; M_3) = \text{lcb}_{j+1}(\underline{y}; M_1)$ for all $0 \leq j \leq 2$.*
- (2) *If \underline{y} is a regular sequence on M_3 then $\text{lcb}_j(\underline{y}; M_1) = \text{lcb}_j(\underline{y}; M_2)$ for all $0 \leq j \leq 3$.*

Proof. The proofs of (1) and (2) are similar and for the sake of brevity we only provide the proof of (1). The sequence \underline{y} is a regular sequence on M_2 and so $H^j(\underline{x}^t; M_2) = 0$ whenever $j \leq 2$. Therefore if $0 \leq j \leq 1$ there are commutative diagrams

$$\begin{array}{ccc} H^j(\underline{y}^t; M_3) & \xrightarrow{\cong} & H^{j+1}(\underline{y}^t; M_1) \\ \downarrow \alpha_{M_3; \underline{y}; t; t+k}^{j+1} & & \downarrow \alpha_{M_1; \underline{y}; t; t+k}^j \\ H^j(\underline{y}^{t+k}; M_3) & \xrightarrow{\cong} & H^{j+1}(\underline{y}^{t+k}; M_1) \end{array}$$

whose horizontal arrows are isomorphisms. It easily follows that $\text{lcb}_j(\underline{y}; M_3) = \text{lcb}_{j+1}(\underline{y}; M_1)$ whenever $0 \leq j \leq 1$. To verify that $\text{lcb}_2(\underline{y}; M_3) = \text{lcb}_3(\underline{y}; M_1)$ consider the following commutative diagrams:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^2(\underline{y}^t; M_3) & \xrightarrow{\delta_t} & H^3(\underline{y}^t; M_1) & \xrightarrow{i_t} & H^3(\underline{y}^t; M_2) \\ & & \downarrow \alpha_{M_3; \underline{y}; t; t+k}^2 & & \downarrow \alpha_{M_1; \underline{y}; t; t+k}^3 & & \downarrow \alpha_{M_2; \underline{y}; t; t+k}^3 \\ 0 & \longrightarrow & H^2(\underline{y}^{t+k}; M_3) & \xrightarrow{\delta_{t+k}} & H^3(\underline{y}^{t+k}; M_1) & \xrightarrow{i_{t+k}} & H^3(\underline{y}^{t+k}; M_2) \end{array}$$

A simple diagram chase and utilizing the injectivity of the maps δ_t, δ_{t+k} , and $\alpha_{M_2; \underline{y}; t; t+k}^3$ imply that $\text{lcb}_2(\underline{y}; M_3) = \text{lcb}_3(\underline{y}; M_1)$. \square

3. ANNIHILATION OF EXT-MODULES AND BOUNDED LOCAL COHOMOLOGY BOUNDS

We are focused on the weak implies strong conjecture and every weakly F -regular ring is a normal Cohen-Macaulay domain, [HH90, Theorem 4.9 and Lemma 5.9]. Therefore we assume throughout this section, and next, that (R, \mathfrak{m}, k) is a normal Cohen-Macaulay domain. We further assume that R is the homomorphic image of a regular local ring S . We write $R \cong S/I$ and h will always denote the height of I , equivalently the codimension of R . In particular, the dimension of S is $h + 4$.

3.1. Annihilation of Ext-modules. The following proposition is our initial step toward providing linearly bounded local cohomology bounds needed to prove Theorem A.

Proposition 3.1. *Let (R, \mathfrak{m}, k) be an excellent local normal Cohen-Macaulay domain of Krull dimension 4. Let $K \subseteq R$ be an ideal of pure height 1 of R so that*

- (1) *The inclusion of ideals $K^i \subseteq K^{(i)}$ is an equality on the punctured spectrum of R ;*
- (2) *The analytic spread of KR_P is no more than $\text{ht}(P) - 1$ for all non-maximal prime ideals P of height at least 2;*

- (3) For each non-maximal prime ideal P of $\text{Spec}(R)$ the ideal KR_P has reduction number 1 with respect to any reduction.

Then there exists an \mathfrak{m} -primary ideal \mathfrak{a} so that

$$\mathfrak{a}^i \text{Ext}_S^{h+3}(R/K^i, S) = 0$$

for all $i \in \mathbb{N}$.

Proof. Choose an element $x_1 \in K$. The principal locus of K is an open subset of $\text{Spec}(R)$ and K localizes to a principal ideal at all height 2 primes of R by assumption. Indeed, an ideal of a normal domain has analytic spread 1 if and only if the ideal is principal. In particular, we can choose a parameter sequence x_2, x_3 of R/x_1R so that KR_{x_2} and KR_{x_3} are principal ideals in their respective localizations. Even further, we replace x_2 and x_3 by x_2^t and x_3^t respectively and may assume that $x_2K \subseteq r_2R$ and $x_3K \subseteq r_3R$ for some elements $r_2, r_3 \in K$. Consider the short exact sequences

$$0 \rightarrow \frac{K^i}{(r_2^i)} \rightarrow \frac{R}{(r_2^i)} \rightarrow \frac{R}{K^i} \rightarrow 0$$

and

$$0 \rightarrow \frac{K^i}{(r_3^i)} \rightarrow \frac{R}{(r_3^i)} \rightarrow \frac{R}{K^i} \rightarrow 0.$$

Observe that x_2^i annihilates $K^i/(r_2^i)$ and x_3^i annihilates $K^i/(r_3^i)$. Therefore

$$\text{Ext}_S^{h+3}(R/K^i, S) \cong \text{Ext}_S^{h+2}(K^i/(r_2^i), S) \cong \text{Ext}_S^{h+2}(K^i/(r_3^i), S)$$

and we find that the ideal (x_1^i, x_2^i, x_3^i) annihilates $\text{Ext}_S^{h+3}(R/K^i, S)$. To complete the proof of the theorem we aim to find a parameter element x_4 of $R/(x_1, x_2, x_3)$ so that x_4^i annihilates $\text{Ext}_S^{h+3}(R/K^i, S)$ for all i .

Let $\Lambda = \{P_1, \dots, P_m\}$ be the prime components of the height 3 parameter ideal (x_1, x_2, x_3) . If necessary, enlarge Λ so that every component of K is contained in some prime ideal of Λ and let $W = R \setminus \bigcup_{P \in \Lambda} P$.

Claim 3.2. *There exist elements $a, b \in K$ such that*

- (1) $(a, b)R_W$ forms a reduction of KR_W ;
- (2) the element a generates K at its components;
- (3) if K' is the unique ideal of pure height 1 whose components are disjoint from K and is such that $(a) = K \cap K'$ then b avoids all components of K' .

Proof of Claim 3.2. We are assuming the ideal K has analytic spread at most 2 at each of the localizations R_P as P varies among the prime ideals in Λ . So for each $1 \leq i \leq m$ there exists $a_i, b_i \in K$ such that $(a_i, b_i)R_{P_i}$ forms a reduction of KR_{P_i} . For each $1 \leq i \leq m$ choose $r_i \in \bigcap_{P \in \Lambda - \{P_i\}} P - P_i$ and set $a' = \sum r_i a_i$ and $b' = \sum r_i b_i$. We claim $(a', b')R_{W_\ell}$ is a reduction of KR_{W_ℓ} . By [HS06, Proposition 8.1.1] it is enough to check (a', b') forms a reduction of K at each of the localizations R_{P_i} for $1 \leq i \leq m$. By [HS06, Proposition 8.2.4] it is enough to check that the fiber cone $R_P/PR_P \otimes R[Kt] \cong \bigoplus K^n R_{P_i}/P_i K^n R_{P_i}$ is finite over the subalgebra spanned by $((a', b')R_{P_i}, P_i K)/P_i K$. But $a' \equiv r_i a_i \pmod{P_i K}$, $b' \equiv r_i b_i \pmod{P_i K}$, r_i is a unit of R_{P_i} , and therefore $(a', b')R_{W_\ell}$ does indeed form a reduction of KR_{W_ℓ} by a second application of [HS06, Proposition 8.2.4].

Now consider the set of primes $\Gamma = \{Q_1, \dots, Q_n\}$ which are the minimal components of K . The purpose of enlarging the set of height ℓ primes in the statement of the claim was to insure that each $Q_j \in \Gamma$ is a prime ideal of the localization R_{W_ℓ} . In particular, $(a', b')R_{Q_i}$ forms a reduction of KR_{Q_i} for each $1 \leq i \leq n$. But R_{Q_i} is a discrete valuation ring and therefore for each $1 \leq i \leq \ell$ either $KR_{Q_i} = (a')R_{Q_i}$ or $KR_{Q_i} = (b')R_{Q_i}$. Without loss of generality we assume that $KR_{Q_i} = (a')R_{Q_i}$ for at least one value of i and relabel the primes in Γ so that $KR_{Q_i} = (a')R_{Q_i}$ for each $1 \leq i \leq j$ and $KR_{Q_i} \neq (a')R_{Q_i}$ for each $j+1 \leq i \leq n$. Choose $r \in Q_1 \cap \dots \cap Q_j - \bigcup_{i=j+1}^n Q_i$ and consider the element $a' + rb'$. We claim that $a' + rb'$ generates KR_{Q_i} for each $1 \leq i \leq n$. First consider a localization at a prime $Q_i \in \Gamma$ with $1 \leq i \leq j$. Then $(a', b')R_{Q_i} = (a')R_{Q_i}$ by assumption and so $(b')R_{Q_i} \subseteq (a')R_{Q_i}$. Because $r \in Q_i$ there is a strict containment of principal ideal $(rb')R_{Q_i} \subsetneq (a')R_{Q_i}$ and it follows that $(a')R_{Q_i} = (a' + rb')R_{Q_i}$. Now consider a localization R_{Q_i} with $j+1 \leq i \leq n$. We are assuming that a' does not generate KR_{Q_i} and therefore $(a')R_{Q_i} \subsetneq (b')R_{Q_i} = KR_{Q_i}$. Moreover, r is a unit of R_{Q_i} and therefore $(b')R_{Q_i} = (a' + rb')R_{Q_i}$.

Let $a = a' + rb'$. Then $(a, b')R_{W_\ell} = (a', b')R_{W_\ell}$ forms a reduction of KR_{W_ℓ} and the element a generates K at each of its minimal components as desired. Suppose as an ideal of R the principal ideal (a) has decomposition $(a) = K \cap K' \cap K''$ so that

- (1) K, K', K'' are pure height 1 ideals whose components are disjoint from one another;
- (2) the components of K' are height 1 prime ideals which do not contain b ;
- (3) the components of K'' are height 1 prime ideals which do contain b .

We take K' or K'' to be R if no such components of (a) exist. If $K'' = R$ then we let $b = b'$ and the elements a, b satisfy the conclusions of the claim. If $K'' \neq R$ then first observe that, because $(a, b')R_{W_\ell}$ forms a reduction of KR_{W_ℓ} and $a, b' \in K''$, we must have that $(a)R_{W_\ell} = (K \cap K')R_{W_\ell}$. Choose an element $r \in K \cap K'$ which avoids all components in K'' and consider the element $b = b' + r$. Then $(a, b)R_{W_\ell} = (a, b')R_{W_\ell}$ forms a reduction of KR_{W_ℓ} . Moreover, the element b avoids all minimal components of K' and K'' by construction. \square

We are assuming that KR_W has reduction number 1 with respect to the reduction $(a, b)R_W$ provided above, i.e. $(a, b)KR_W = K^2R_W$. The following R -modules localize to 0 over R_W :

- (1) $K^2/(a, b)K$;
- (2) $\text{Ext}_S^{h+3}(R/(a, b)K, S)$;
- (3) $\text{Ext}_S^{h+3}(R/K, S)$.

Therefore there exists a parameter element x_4 of $R/(x_1, x_2, x_3)$ so that

- (1) $x_4K^2 \subseteq (a, b)K$;
- (2) $x_4 \text{Ext}_S^{h+3}(R/(a, b)K, S) = 0$;
- (3) $x_4 \text{Ext}_S^{h+3}(R/K, S) = 0$.

Even further, as $x_4K^2 \subseteq (a, b)K$, observe that $x_4^{i-1}K^i \subseteq (a, b)^{i-1}K$ for all $i \in \mathbb{N}$. Consider the following short exact sequence:

$$0 \rightarrow \frac{K^i}{(a, b)^{i-1}K} \rightarrow \frac{R}{(a, b)^{i-1}K} \rightarrow \frac{R}{K^i} \rightarrow 0.$$

The left most term is annihilated by x_4^{i-1} . Thus, in order to show x_4^i annihilates $\text{Ext}_S^{h+3}(R/K^i, S)$ it suffices to show that $x_4 \in \bigcap_{i \in \mathbb{N}} \text{Ann}_R(\text{Ext}_S^{h+3}(R/(a, b)^{i-1}K, S))$. To this end we present a claim:

Claim 3.3. *For every integer i there is a short exact sequence*

$$0 \rightarrow \frac{R}{ab^i R} \rightarrow \frac{R}{a(a, b)^{i-1}K} \oplus \frac{R}{b^i K} \rightarrow \frac{R}{(a, b)^i K} \rightarrow 0.$$

Proof of Claim 3.3. For any ideals I, J there is a short exact sequence

$$0 \rightarrow \frac{R}{I \cap J} \rightarrow \frac{R}{I} \oplus \frac{R}{J} \rightarrow \frac{R}{I + J} \rightarrow 0.$$

Thus to prove the claim we need only to observe $a(a, b)^{i-1}K \cap b^i K = ab^i R$. Clearly $ab^i \in a(a, b)^{i-1}K \cap b^i K$. On the other hand, an element of $a(a, b)^{i-1}K \cap b^i K$ is of the form $b^i r$ where $r \in K$ and $b^i r \in a(a, b)^{i-1}K$. To show that $b^i r \in ab^i R$ we must show that $r \in aR = K \cap K'$. The element $r \in K$ by assumption. Localizing at a component P of K' , a component which does not contain b by design, we find that $rb^i \in a(a, b)^{i-1}R_P = aR_P$ and thus $r \in aR$ as desired. \square

Claim 3.3 provides to us isomorphisms

$$(3.1) \quad \text{Ext}_S^{h+3}(R/(a, b)^i K, S) \cong \text{Ext}_S^{h+3}(R/a(a, b)^{i-1}K, S) \oplus \text{Ext}_S^{h+3}(R/b^i K, S).$$

There are short exact sequences

$$0 \rightarrow \frac{R}{(a, b)^{i-1}K} \xrightarrow{\cdot a} \frac{R}{a(a, b)^{i-1}K} \rightarrow \frac{R}{aR} \rightarrow 0$$

and

$$0 \rightarrow \frac{R}{K} \xrightarrow{\cdot b^i} \frac{R}{b^i K} \rightarrow \frac{R}{b^i R} \rightarrow 0.$$

Therefore the isomorphisms of (3.1) can be further rewritten as

$$\text{Ext}_S^{h+3}(R/(a, b)^i K, S) \cong \text{Ext}_S^{h+3}(R/(a, b)^{i-1}K, S) \oplus \text{Ext}_S^{h+3}(R/K, S).$$

Inductively, we find that

$$\text{Ext}_S^{h+3}(R/(a, b)^i K, S) \cong \bigoplus^i \text{Ext}_S^{h+3}(R/K, S)$$

and we conclude that x_4 annihilates $\text{Ext}_S^{h+3}(R/(a, b)^i K, S)$ as desired. \square

Suppose that $I \subseteq R$ is an unmixed ideal. The Rees algebra of I is the standard graded R -algebra $R[It] = \bigoplus_{N \geq 0} I^N$, the associated graded ring of I is $\text{Gr}_I(R) = R[It] \otimes_R R/I$, and the symbolic Rees algebra of I is $\mathcal{R}_I = \bigoplus_{N \geq 0} I^{(N)}$. The inclusion of \mathbb{N} -graded R -algebras $R[It] \subseteq \mathcal{R}_I$ is an equality if and only if $\bigcup_{N \in \mathbb{N}} \text{Ass}(R/I^N)$ agrees with the set of minimal prime ideals of I . The R -algebra \mathcal{R}_I is Noetherian if and only if there exists an m so that $\mathcal{R}_{I^{(m)}}$ is standard graded, i.e. $I^{(m)N} = I^{(mN)}$ for all N .

Suppose that \mathcal{R}_I is Noetherian and m is chosen so that $\mathcal{R}_{I^{(m)}}$ is standard graded. Let $\ell(I^{(m)}) = \dim \mathcal{R}_{I^{(m)}} \otimes_R R/\mathfrak{m}$, the analytic spread of $I^{(m)}$. If m' is another integer so that $\mathcal{R}_{I^{(m')}}$ is standard graded then $I^{(m)m'} = I^{(m')m} = I^{(mm')}$ and so $\ell(I^{(m)}) = \ell(I^{(mm')}) = \ell(I^{(m)})$.

We say that *the associated graded rings of \mathcal{R}_I have negative a -invariants* if the graded local cohomology modules $H_{\mathcal{R}_I^+}^\ell(\mathrm{Gr}_{I^{(m)}}(R))$ are supported only in negative degrees where $\ell = \ell(I^{(m)})$ is the analytic spread of a choice of symbolic power of I for which $R[I^{(m)}t] = \mathcal{R}_{I^{(m)}}$.

Suppose that R is a normal domain, $J_1 \subset R$ is a choice of canonical ideal, and $x_1 \in J_1$ is a choice of generic generator. Then we can write $(x_1) = J_1 \cap K_1$ where K_1 is an ideal of pure height 1 whose components are disjoint from J_1 . Then K_1 is an *anticanonical ideal* of R and we refer to the symbolic Rees algebra \mathcal{R}_{K_1} as the *anticanonical algebra* of R .

Theorem 3.4. *Let (R, \mathfrak{m}, k) be an excellent local normal Cohen-Macaulay domain of Krull dimension 4. Suppose that the anticanonical algebra of R is Noetherian on the punctured spectrum of R so that its associated graded rings have negative a -invariant. Let $J_1 \subseteq R$ be a choice of canonical ideal of R . Then there exists an integer $m \in \mathbb{N}$ and \mathfrak{m} -primary ideal $\mathfrak{a} \subseteq R$ so that $J_1^{(m)}$ is principal in codimension 2 and*

$$\mathfrak{a}^i \mathrm{Ext}_S^{h+2}(\mathrm{Ext}_S^{h+1}(R/J_1^{mi+1}, S), S) = 0$$

for every integer $i \in \mathbb{N}$.

Proof. Choose a generic generator $x_1 \in J_1$ and write $(x_1) = J_1 \cap K_1$ where K_1 is an anticanonical ideal of R whose components are disjoint from the components of J_1 . A theorem of Brodmann, [Bro79], asserts that $\Gamma := \bigcup^\infty \mathrm{Ass}(R/K_1^n)$ is a finite set, c.f. [HS15]. Let $P_1, \dots, P_t \in \Gamma$ be the finitely many non-maximal primes of Γ which are not of height 1. If P is a non-maximal ideal of R not belonging to $\{P_1, \dots, P_t\}$ then $K_1^i R_P = K_1^{(i)} R_P$ for all i . If $P_j \in \{P_1, \dots, P_t\}$ then our assumption that the anticanonical algebra is Noetherian on the punctured spectrum implies that there exists an integer m_j so that $K_1^{(m_j)i} R_P = K_1^{(m_j i)} R_P$ and that the analytic spread of $K_1^{(m_j i)} R_P$ does not exceed $\mathrm{ht}(P) - 1$, see [KR86] and [CHS10, Theorem 1.5].

Let m be a common multiple of m_1, \dots, m_t . Then the inclusion of ideals $K_1^{(m)i} \subseteq K_1^{(mi)}$ becomes an equality when localized at any non-maximal prime ideal of R . The ideal $K_1^{(m)}$ is principal in codimension 2 since its analytic spread in codimension 2 is 1. As $J_1^{(m)}$ is the inverse element of $K_1^{(m)}$ when viewed as elements of the divisor class group of R , $J_1^{(m)}$ is principal in codimension 2 as well. Even further, because we are assuming the a -invariant of the associated graded ring of the anticanonical algebra of R_P is negative for each non-maximal prime ideal R , we can replace m by a multiple of itself and assume that the ideal $K_1^{(m)} R_{P_j}$ has reduction number 1 with respect to any reduction, see [Hoa93, Theorem 2.1]. Let $K = K_1^{(m)}$ and $x = x_1^m$.

Claim 3.5. *For each integer i*

$$\mathrm{Ext}_S^{h+2}(\mathrm{Ext}_S^{h+1}(R/J_1^{mi+1}, S), S) \cong \mathrm{Ext}_S^{h+3}(R/K^i, S).$$

Proof of Claim 3.5. For each integer i there is short exact sequence

$$0 \rightarrow \frac{J_1^{mi+1}}{x_1^{mi+1} J_1} \rightarrow \frac{R}{x_1^{mi+1} J_1} \rightarrow \frac{R}{J_1^{mi+1}} \rightarrow 0.$$

The ideal $x_1^{mi+1}J_1$ is a canonical ideal of R . Therefore

$$\mathrm{Ext}_S^{h+1}(R/x_1^{mi+1}, S) \cong R/x_1^{mi+1}J_1$$

and there are left exact sequences

$$0 \rightarrow \mathrm{Ext}_S^{h+1}(R/J_1^{mi+1}, S) \rightarrow \frac{R}{x_1^{mi+1}J_1} \rightarrow \mathrm{Ext}_S^{h+1}(J_1^{mi+1}/x_1^{mi+1}J_1, S).$$

Therefore $\mathrm{Ext}_S^{h+1}(R/J_1^{mi+1}, S) \cong L_i/x_1^{mi+1}J_1$ for some ideal $L_i \subseteq R$. Moreover, $R/L_i \subseteq \mathrm{Ext}_S^{h+1}(J_1^{mi+1}/x_1^{mi+1}J_1, S)$. Because $\mathrm{Ext}_S^{h+1}(J_1^{mi+1}/x_1^{mi+1}J_1, S)$ is an (S_2) -module over its support it follows that R/L_i is an (S_1) -module over its support. Hence L_i , as an ideal of R , is unmixed of height 1. Moreover, every component of L_i is a component of x_1R . Localizing at a component of J_1 we see that L_i agrees with x_1R and localizing at a component of K_1 we see that L_i agrees with x_1^{mi+1} . Therefore L_i agrees with the unmixed ideal $x_1K_1^{(mi)}$ and so

$$\mathrm{Ext}_S^{h+1}(R/J_1^{mi+1}, S) \cong x_1K_1^{(mi)}/x_1^{mi+1}J_1.$$

If we divide by x_1 we find that

$$x_1K_1^{(mi)}/x_1^{mi+1}J_1 \cong K_1^{(mi)}/x_1^{mi}J_1 = K^{(i)}/x^iJ_1.$$

Now we consider the short exact sequences

$$0 \rightarrow K^{(i)}/x^iJ_1 \rightarrow R/x^iJ_1 \rightarrow R/K^{(i)} \rightarrow 0.$$

The cyclic R -module R/x^iJ_1 is Cohen-Macaulay of dimension 3 and therefore

$$\mathrm{Ext}_S^{h+2}(K^{(i)}/x^iJ_1, S) \cong \mathrm{Ext}_S^{h+3}(R/K^{(i)}, S).$$

To complete the proof of the claim we consider the short exact sequences

$$0 \rightarrow \frac{K^{(i)}}{K^i} \rightarrow \frac{R}{K^i} \rightarrow \frac{R}{K^{(i)}} \rightarrow 0.$$

The inclusion of ideals $K^i \subseteq K^{(i)}$ are an equality on the punctured spectrum of R and hence

$$\mathrm{Ext}_S^{h+3}(R/K^{(i)}, S) \cong \mathrm{Ext}_S^{h+3}(R/K^i, S).$$

□

To prove the theorem we aim to find an \mathfrak{m} -primary ideal so that

$$\mathfrak{a}^i \mathrm{Ext}_S^{h+3}(R/K^i, S) = 0$$

for all i . Such an annihilation property is the content of Proposition 3.1.

□

3.2. Existence of bounded local cohomology bounds. Let $I \subseteq R$ be an ideal of pure height 1 and $(F_\bullet, \partial_\bullet)$ be an S -free resolution of R/I and G_\bullet an S -free resolution of $\omega_{R/I} \cong \text{Ext}_S^{h+1}(R/I, S)$. Let $(-)^* = \text{Hom}_S(-, S)$. Then $H^i(F_\bullet^*) = \text{Ext}_S^i(R/I, S) = 0$ for all $i \leq h$ and if we let \tilde{F}_\bullet^* be the truncation of F_\bullet^* at the $h+1$ st spot then \tilde{F}_\bullet^* resolves $\text{Coker}(\partial_{h+1}^*)$. There is a natural inclusion of S -modules $\text{Ext}_S^{h+1}(R/I, S) \subseteq \text{Coker}(\partial_{h+1}^*)$. This inclusion produces a map of complex to cocomplex $G_\bullet \rightarrow \tilde{F}_\bullet^*$ and thus a natural map $R/I \rightarrow \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/I, S), S)$ obtained from applying $\text{Hom}_S(-, S)$ to $G_\bullet \rightarrow \tilde{F}_\bullet^*$. The map $R/I \rightarrow \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/I, S), S)$ is injective and is an isomorphism in the Cohen-Macaulay locus of R/I .

Theorem 3.6. *Let (R, \mathfrak{m}, k) be an excellent local normal Cohen-Macaulay domain of Krull dimension 4. Suppose that the anticanonical algebra of R is Noetherian on the punctured spectrum of R so that its associated graded rings have negative a -invariant. Let $J_1 \subseteq R$ be a choice of canonical ideal of R and \mathfrak{a} as in Theorem 3.4. There exists an integer m and $x_1 \in J_1$, such that if $x_2, x_3, x_4 \in \mathfrak{a}$ are parameters on R/x_1R chosen so that*

- (1) $x_2J_1 \subseteq a_2R$ for some $a_2 \in J_1$;
- (2) $x_3J_1^{(m)} \subseteq a_3R$ for some $a_3 \in J_1^{(m)}$.

Then for each natural number i there exists an integer ℓ such that

$$\text{lcb}_3(x_2^\ell, x_3^\ell, x_4; R/J_1^{(mi+1)}) \leq i + 1.$$

Proof. Let m and \mathfrak{a} be as in Theorem 3.4. Because J_1 is principal in codimension 1 and $J_1^{(m)}$ is principal in codimension 2 we can choose a parameter sequence $x_2, x_3, x_4 \in \mathfrak{a}$ on R/x_1R so that

- (1) $x_2J_1 \subseteq a_2R$ for some $a_2 \in J_1$;
- (2) $x_3J_1^{(m)} \subseteq a_3R$ for some $a_3 \in J_1^{(m)}$.

For each integer i there is a short exact sequence of the form

$$0 \rightarrow R/J_1^{(mi+1)} \rightarrow \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/J_1^{(mi+1)}, S), S) \rightarrow C_i \rightarrow 0.$$

Inverting the element x_2 or x_3 has the effect of making the ideal J_1 principal. Therefore the first map in the above short exact sequence is an isomorphism whenever x_2 or x_3 is inverted and so there exists an integer ℓ so that x_2^ℓ, x_3^ℓ annihilates C_i . By (3) of Proposition 2.4 we have that

$$\text{lcb}_3(x_2^\ell, x_3^\ell, x_4; R/J_1^{(mi+1)}) \leq \text{lcb}_3(x_2^\ell, x_3^\ell, x_4; \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/J_1^{(mi+1)}, S), S)) + 1.$$

Our aim is to show $\text{lcb}_3(x_2^\ell, x_3^\ell, x_4; \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/J_1^{(mi+1)}, S), S)) \leq i$. By Lemma 2.2 it suffices to prove $\text{lcb}_3(x_2^{\ell i}, x_3^{\ell i}, x_4^i; \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/J_1^{(mi+1)}, S), S)) \leq 1$.

Let $(F_\bullet, \partial_\bullet)$ be the minimal free S -resolution of $\text{Ext}_S^{h+1}(R/J^{(mi+1)}, S)$ and let $(-)^* = \text{Hom}_R(-, S)$. The module $\text{Ext}_S^{h+1}(R/J^{(mi+1)}, S)$ has depth at least 2 and so $F_{h+3} = F_{h+4} = 0$. It follows that there are short exact sequences

$$0 \rightarrow \text{Im}(\partial_{h+2}^*) \rightarrow F_{h+2}^* \rightarrow \text{Ext}_S^{h+2}(\text{Ext}_S^{h+1}(R/J^{mi+1}, S), S) \rightarrow 0$$

and

$$0 \rightarrow \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/J^{mi+1}, S), S) \rightarrow \text{Coker}(\partial_{h+1}^*) \rightarrow \text{Im}(\partial_{h+2}^*) \rightarrow 0.$$

The S -module $\text{Coker}(\partial_{h+1}^*)$ has projective dimension $h + 1$ and is annihilated by the height $h + 1$ ideal J^{mi+1} . By a simple prime avoidance argument we may lift x_2, x_3, x_4 to elements of S so that x_2, x_3, x_4 forms a regular sequence on $\text{Coker}(\partial_{h+1}^*)$ and the free S -module F_{h+2}^* . By two applications of (1) of Proposition 2.5 applied to the two above short exact sequences

$$\begin{aligned} \text{lcb}_3(x_2^{\ell i}, x_3^{\ell i}, x_4^i; \text{Ext}_S^{h+1}(\text{Ext}_S^{h+1}(R/J_1^{mi+1}, S), S)) &= \text{lcb}_2(x_2^{\ell i}, x_3^{\ell i}, x_4^i; \text{Im}(\partial_{h+2}^*)) \\ &= \text{lcb}_1(x_2^{\ell i}, x_3^{\ell i}, x_4^i; \text{Ext}_S^{h+2}(\text{Ext}_S^{h+1}(R/J_1^{mi+1}, S), S)). \end{aligned}$$

By Theorem 3.4 we have that

$$(x_2^{\ell i}, x_3^{\ell i}, x_4^i) \text{Ext}_S^{h+2}(\text{Ext}_S^{h+1}(R/J_1^{mi+1}, S), S) = 0.$$

Therefore

$$\text{lcb}_1(x_2^{\ell i}, x_3^{\ell i}, x_4^i; \text{Ext}_S^{h+2}(\text{Ext}_S^{h+1}(R/J_1^{mi+1}, S), S)) = 1$$

by (1) of Proposition 2.3. □

4. EQUALITY OF TEST IDEALS

If $N \subseteq M$ are R -modules then the *finitistic tight closure of N inside M* is the union of $(N \cap M')_{M'}^*$ where $M' \subseteq M$ runs through all finitely generated submodules of M . Let $E_R(k)$ be the injective hull of the residue field of (R, \mathfrak{m}, k) . Then R is strongly F -regular if and only if $0 = 0_{E_R(k)}^*$, [HH90, Proposition 8.23] and R is weakly F -regular if and only if $0_{E_R(k)}^{*,fg} = 0$, [Smi93, Proposition 7.1.2].

With the exception of Lemma 4.3 and Corollary 4.5, we continue to assume that (R, \mathfrak{m}, k) is an excellent normal Cohen-Macaulay domain of Krull dimension 4 and is the homomorphic image of a regular local ring S so that the results of Section 3 are applicable. We fix the characteristic of R to be of prime characteristic $p > 0$.

The following lemma is inspired by the methodology of Williams and MacCrimmon, [Wil95, Mac96]. The lemma is well-known by experts, can be pieced together by work of the first author in [Abe02], and we refer the reader to [PT18, Lemma 6.7] for a more general statement.

Lemma 4.1. *Suppose that (R, \mathfrak{m}, k) is a local normal Cohen-Macaulay domain of prime characteristic $p > 0$, of Krull dimension 4, and $J \subseteq R$ an ideal of pure height 1. Let $y_1 \in J$ and y_2, y_3, y_4 parameters on R/y_1R and fix $e \in \mathbb{N}$.*

- (1) *If $y_2J \subseteq aR$ for some $a \in J$, then for any integers N_2, N_3, N_4 with $N_2 \geq 2$, we have that*

$$\begin{aligned} & ((J^{(p^e)}, y_2^{N_2 p^e}, y_3^{N_3 p^e}, y_4^{N_4 p^e}) : y_2^{(N_2-1)p^e}) \\ &= ((J^{[p^e]}, y_2^{N_2 p^e}, y_3^{N_3 p^e}, y_4^{N_4 p^e}) : y_2^{(N_2-1)p^e}) \\ &= ((J^{[p^e]}, y_2^{2p^e}, y_3^{N_3 p^e}, y_4^{N_4 p^e}) : y_2^{p^e}). \end{aligned}$$

- (2) *Suppose $y_3J^{(m)} \subseteq bR$ for some $b \in J^{(m)}$, then for any non-negative integers N_2, N_3, N_4 with $N_3 \geq 2$, we have that*

$$\begin{aligned} & ((J^{(p^e)}, y_2^{N_2 p^e}, y_3^{N_3 p^e}, y_4^{N_4 p^e}) : y_3^{(N_3-1)p^e}) \\ & \subseteq ((J^{(p^e)}, y_2^{N_2 p^e}, y_3^{2p^e}, y_4^{N_4 p^e}) : y_1^m y_3^{p^e}). \end{aligned}$$

Theorem 4.2. *Let (R, \mathfrak{m}, k) be an excellent local normal Cohen-Macaulay domain of prime characteristic $p > 0$ and of Krull dimension 4. Suppose that the anticanonical algebra of R is Noetherian on the punctured spectrum of R so that its associated graded rings have negative a -invariant. Then*

$$0_{E_R(k)}^{*,fg} = 0_{E_R(k)}^*.$$

In particular, if R is weakly F -regular then R is strongly F -regular.

Proof. Let $J_1 \subset R$ be a canonical ideal of R and let x_1, x_2, x_3, x_4 and m be as in the statement of Theorem 3.6. Identify the injective hull $E_R(k)$ as

$$E_R(k) = \varinjlim \left(\frac{R}{(x_1^{t-1}J_1, x_2^t, x_3^t, x_4^t)} \xrightarrow{\cdot x_1 x_2 x_3 x_4} \frac{R}{(x_1^t J_1, x_2^{t+1}, x_3^{t+1}, x_4^{t+1})} \right).$$

Suppose that $\eta = [r + (x_1^{t-1}J_1, x_2^t, x_3^t, x_4^t)] \in 0_{E_R(k)}^*$. Equivalently, if $c \in R$ is a test element then for all $e \in \mathbb{N}$ there exists an integer s so that

$$cr^{p^e} (x_1 x_2 x_3 x_4)^{sp^e} \in (x_1^{t+s-1} J_1, x_2^{t+s}, x_3^{t+s}, x_4^{t+s})^{[p^e]}.$$

The element x_1 is regular on $R/(x_2, x_3, x_4)$ and therefore

$$cr^{p^e} (x_2 x_3 x_4)^{sp^e} \in (x_1^{t-1} J_1, x_2^{t+s}, x_3^{t+s}, x_4^{t+s})^{[p^e]} = (J, x_2^{t+s}, x_3^{t+s}, x_4^{t+s})^{[p^e]}.$$

where $J = x_1^{t-1} J_1$. Multiplying by $(x_2 x_3 x_4)^{p^e(t-s-(t+s))}$ we find that

$$cr^{p^e} (x_2^t x_3^t x_4^t)^{(s-1)p^e} \in (J, x_2^{ts}, x_3^{ts}, x_4^{ts})^{[p^e]}.$$

Let y_2, y_3, y_4 denote the parameter sequence x_2^t, x_3^t, x_4^t so that

$$cr^{p^e} (y_2 y_3 y_4)^{(s-1)p^e} \in (J, y_2^s, y_3^s, y_4^s)^{[p^e]}.$$

Observe that $p^e \geq m(\lfloor \frac{p^e}{m} \rfloor - 1) + 1$ and hence $J^{[p^e]} \subseteq J^{(p^e)} \subseteq J^{(mi+1)}$ where we set $i = \lfloor \frac{p^e}{m} \rfloor - 1$. Therefore

$$(4.1) \quad cr^{p^e} (y_2 y_3 y_4)^{(s-1)p^e} \in (J^{(mi+1)}, y_2^{sp^e}, y_3^{sp^e}, y_4^{sp^e}).$$

Let ℓ be the integer depending on i described in Theorem 3.6. Theorem 3.6 and Lemma 2.2 tell us that for each integer i that there exists an integer ℓ so that

$$\text{lcb}_3(y_2^\ell, y_3^\ell, y_4; R/J_1^{(mi+1)}) \leq i + 1.$$

Because $J = x_1^{t-1} J_1$ we have that for each integer i there is a short exact sequence

$$0 \rightarrow \frac{R}{J_1^{(mi+1)}} \xrightarrow{\cdot x_1^{(t-1)(mi+1)}} \frac{R}{J^{(mi+1)}} \rightarrow \frac{R}{x_1^{(t-1)(mi+1)} R} \rightarrow 0.$$

The sequence y_2^ℓ, y_3^ℓ, y_4 is a regular sequence on $R/x_1^{(t-1)(mi+1)} R$. By (2) of Proposition 2.5 we have that

$$(4.2) \quad \text{lcb}_3(y_2^\ell, y_3^\ell, y_4; R/J_1^{(mi+1)}) = \text{lcb}_3(y_2^\ell, y_3^\ell, y_4; R/x_1^{t-1} J_1^{(mi+1)}) \leq i + 1.$$

We multiply the containment (4.1) by $(y_2 y_3)^{(\ell-1)sp^e}$ and notice that

$$(4.3) \quad \begin{aligned} cr^{p^e} (y_2 y_3 y_4)^{(s-1)p^e} (y_2 y_3)^{(\ell-1)sp^e} &= cr^{p^e} (y_2 y_3)^{(\ell-1)p^e} (y_2^\ell y_3^\ell y_4)^{(s-1)p^e} \\ &\in (J^{(mi+1)}, y_2^{\ell sp^e}, y_3^{\ell sp^e}, y_4^{sp^e}). \end{aligned}$$

Consider the element

$$\zeta = [cr^{p^e}(y_2y_3)^{(\ell-1)p^e} + (y_2^{\ell p^e}, y_3^{\ell p^e}, y_4^{p^e})]$$

of the top Koszul cohomology group

$$H^3(y_2^{\ell p^e}, y_3^{\ell p^e}, y_4^{p^e}; R/J^{(mi+1)}).$$

Using the notation of Section 2, the containment of (4.3) is equivalent to the assertion that

$$\alpha_{R/J^{(mi+1)}; y_2^{\ell}, y_3^{\ell}, y_4, p^e, sp^e}(\zeta) = 0 \in H^3(y_2^{\ell sp^e}, y_2^{\ell sp^e}, y_4^{sp^e}; R/J_1^{(mi+1)}).$$

By (4.2) we have that

$$\text{lcb}_3(y_2^{\ell}, y_3^{\ell}, y_4; R/J^{(mi+1)}) \leq i + 1 \leq p^e - 1 + 1 = p^e$$

and therefore

$$\alpha_{R/J^{(mi+1)}; y_2^{\ell}, y_3^{\ell}, y_4, p^e, 2p^e}(\zeta) = 0 \in H^3(y_2^{\ell 2p^e}, y_2^{\ell 2p^e}, y_4^{2p^e}; R/J_1^{(mi+1)}).$$

Equivalently, the element

$$cr^{p^e}(y_2y_3)^{(\ell-1)p^e}(y_2^{\ell}y_3^{\ell}y_4)^{p^e} = c(ry_4)^{p^e}y_2^{(2\ell-1)p^e}y_3^{(2\ell-1)p^e} \in (J^{(mi+1)}, y_2^{2\ell p^e}, y_3^{2\ell p^e}, y_4^{2p^e}).$$

Recall that $i = \lfloor \frac{p^e}{m} \rfloor - 1$ and so $m \lfloor \frac{p^e}{m} \rfloor \geq m(\frac{p^e}{m} - 1) = p^e - m$. Hence $mi + 1 \geq p^e - (2m - 1)$ and so

$$c(ry_4)^{p^e}y_2^{(2\ell-1)p^e}y_3^{(2\ell-1)p^e} \in (J^{(p^e-(2m-1))}, y_2^{2\ell p^e}, y_3^{2\ell p^e}, y_4^{2p^e}).$$

Pick a nonzero element $z \in J^{(2m-1)}$. Then

$$zc(ry_4)^{p^e}y_2^{(2\ell-1)p^e}y_3^{(2\ell-1)p^e} \in (J^{(p^e)}, y_2^{2\ell p^e}, y_3^{2\ell p^e}, y_4^{2p^e}).$$

We want to utilize Lemma 4.1 to simplify the above containment. Recall that $J = x_1^{t-1}J_1$, $y_2 = x_2^t$, $y_3 = x_3^t$, $x_2J_1 \subseteq a_2R$ for some $a_2 \in J_1$, and $x_3J_1^{(m)} \subseteq a_3R$ for some $a_3 \in J_1^{(m)}$. Then $y_2J \subseteq x_1^{t-1}a_2R$, $x_1^{t-1}a_2 \in J$, $y_3J^{(m)} \subseteq x_1^{(t-1)m}a_3R$, and $x_1^{(t-1)m}a_3 \in J^{(m)}$. Moreover, $y_1 = x_1^t \in J$ and y_2, y_3, y_4 is a parameter sequence on R/y_1R . Therefore we can apply (2) of Lemma 4.1 and conclude that

$$y_1^m zc(ry_3y_4)^{p^e}y_2^{(2\ell-1)p^e} \in (J^{(p^e)}, y_2^{2\ell p^e}, y_3^{2p^e}, y_4^{2p^e}).$$

By (1) of Lemma 4.1 we are then able to assert that

$$y_1^m zc(ry_2y_3y_4)^{p^e} \in (J^{[p^e]}, y_2^{2p^e}, y_3^{2p^e}, y_4^{2p^e}).$$

The element $y_1^m zc$ does not depend on e . Therefore

$$ry_2y_3y_4 \in (J, y_2^2, y_3^2, y_4^2)^*$$

and hence, as an element of $E_R(k)$, $\eta = [ry_2y_3y_4 + (J, y_2^2, y_3^2, y_4^2)]$ belongs to $0_{E_R(k)}^{*,fg}$. \square

To utilize Theorem 4.2 and prove Theorem A we must observe that the a -invariant of the associated graded rings of the anticanonical algebra is negative whenever the ambient ring is strongly F -regular and the anticanonical algebra is Noetherian.

Lemma 4.3. *Let (R, \mathfrak{m}, k) be an excellent strongly F -regular ring of prime characteristic $p > 0$ and Krull dimension $d \geq 2$. Suppose that $I \subseteq R$ is an ideal of pure height 1 such that $I^N = I^{(N)}$ for all N . Then the associated graded ring of I has negative a -invariant.*

Proof. If $\ell(I) = 1$ then I is principal and $H_{R[It]^+}^1(\mathrm{Gr}_I(R)) = 0$. So we may assume that $2 \leq \ell(I) \leq d$. We first observe that $a_i(R[It]) < 0$ for all $2 \leq i \leq d$. Because $I^N = I^{(N)}$ for all N we have that $S := R[It]$ is a strongly F -regular graded R -algebra by [CEMS18b, Lemma 3.1], see also [Wat94, Theorem 0.1] and [MPST19, Main Theorem]. The cohomology groups $H_{S_+}^i(S)$ are only supported in finitely many positive degrees. Indeed, let $X = \mathrm{Proj}(S)$ so that $H_{S_+}^i(S) \cong H^{i-1}(X, \mathcal{O}_X)$ for all $i \geq 2$, see [ILL+07, Theorem 12.41], and therefore $[H_{S_+}^i(S)]_N = H^{i-1}(X, \mathcal{O}_X(N)) = 0$ for all $N \gg 0$ by Serre vanishing, [Har77, Theorem 5.2]. It follows that there exists a homogeneous positive degree element $c \in S$ such that $c[H_{S_+}^i(S)]_{\geq 0} = 0$. Because S is strongly F -regular the S -linear maps $S \xrightarrow{\cdot F_*^e c} F_*^e S$ are pure for all $e \gg 0$. Therefore the e th Frobenius action on $H_{S_+}^i(S)$ followed by multiplying by c , which is the map realized by tensoring the pure map $S \xrightarrow{\cdot F_*^e c} F_*^e S$ with $H_{S_+}^i(S)$, are injective. But the e th Frobenius action of $H_{S_+}^i(S)$ maps elements of degree n to elements of degree np^e . Furthermore, c was chosen to annihilate elements of non-negative degree and therefore $H_{S_+}^i(S)$ can only be supported in negative degree.

The ring $S = R[It]$ is Cohen-Macaulay and therefore $a_d(\mathrm{gr}_I(R)) < 0$ by [Hoa93, Theorem 3.1]. By [Tru98, Theorem 3.1 (ii)] we have that $a_i(\mathrm{gr}_I(R)) = a_i(S)$ whenever $a_i(\mathrm{gr}_I(R)) \geq a_{i+1}(\mathrm{gr}_I(R))$. An easy descending induction argument now tells us that $a_i(\mathrm{gr}_I(R)) < 0$ for all $2 \leq i \leq d$ and this completes the proof of the theorem. \square

Corollary 4.4. *Let R be an excellent 4-dimensional weakly F -regular ring of prime characteristic $p > 0$. If the anticanonical algebra of R is Noetherian on the punctured spectrum of R then R is strongly F -regular.*

Proof. It is well known that the properties of being weakly F -regular and strongly F -regular can be checked at localizations at the maximal ideals of R , see [HH90, Corollary 4.15]. Thus we may assume $R = (R, \mathfrak{m}, k)$ is local. The properties of weakly F -regular and strongly F -regular for a local ring can be checked after completion. In which case, the property of being weakly F -regular is equivalent to $0_{E_R(k)}^{*,fg}$ being 0 and the property of being strongly F -regular is equivalent to $0_{E_R(k)}^*$ being 0.

Using gamma constructions with respect to a choice of coefficient field, we may assume R is F -finite, see [HH94a, Section 6 and Theorem 7.24] and [Has10, Corollary 3.31]. Every complete local weakly F -regular ring is a normal Cohen-Macaulay domain by [HH90, Lemma 5.9 and Theorem 4.9]. Every weakly F -regular ring is a splinter, [HH94b, Corollary 5.23]. The property of being a splinter localizes. Therefore R is strongly F -regular on the punctured spectrum of R by [CEMS18a, Corollary 5.9]. Lemma 4.3 tells us the anticanonical algebra of R is such that its associated graded ring has negative a -invariant on the punctured spectrum of R and therefore R is strongly F -regular by Theorem 4.2. \square

Corollary 4.5. *Let R be a 4-dimensional normal Cohen-Macaulay domain of prime characteristic p . Suppose that either*

- (1) *R is finitely generated over a field of prime characteristic $p > 5$ with infinite transcendence degree over \mathbb{F}_p and is weakly F -regular;*
- (2) *R is essentially of finite type over a field of prime characteristic $p > 5$ and is F -regular.*

Then R is strongly F -regular.

Proof. Every weakly F -regular ring that is finitely generated over a field of infinite transcendence degree over \mathbb{F}_p is F -regular by [HH94a, Theorem 8.1]. Thus it suffices to prove statement (2) only.

As in the proof of Corollary 4.4, we may assume $R = (R, \mathfrak{m}, k)$ is local and we can use gamma constructions to reduce to the scenario that R is F -finite. The ring R is strongly F -regular at non-maximal points by [Wil95, Main Result]. By [SS10, Corollary 6.9], if P is a nonmaximal prime ideal of R then there exists an effective boundary divisor Δ such that $(\text{Spec}(R_P), \Delta)$ is globally F -regular (or just F -regular since $\text{Spec}(R_P)$ is affine) and therefore has KLT singularities by [HW02, Theorem 3.3]. By [DW19, Corollary 1.12] the anticanonical algebra of R_P is Noetherian and therefore R is strongly F -regular by Corollary 4.4. \square

Acknowledgements

This article would not have been possible without significant feedback from the community. We are most grateful to Craig Huneke, Linqun Ma, and Sarasij Maitra for their considerations and time.

REFERENCES

- [Abe02] Ian M. Aberbach. Some conditions for the equivalence of weak and strong F -regularity. *Comm. Algebra*, 30(4):1635–1651, 2002. [12](#)
- [AP21] Ian Aberbach and Thomas Polstra. Local cohomology bounds and test ideals, 2021. [2](#)
- [BH93] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*, volume 39 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993. [3](#)
- [Bro79] M. Brodmann. Asymptotic stability of $\text{Ass}(M/I^n M)$. *Proc. Amer. Math. Soc.*, 74(1):16–18, 1979. [9](#)
- [CEMS18a] Alberto Chiechio, Florian Enescu, Lance Edward Miller, and Karl Schwede. Test ideals in rings with finitely generated anti-canonical algebras. *J. Inst. Math. Jussieu*, 17(1):171–206, 2018. [15](#)
- [CEMS18b] Alberto Chiechio, Florian Enescu, Lance Edward Miller, and Karl Schwede. Test ideals in rings with finitely generated anti-canonical algebras—corrigendum [MR3742559]. *J. Inst. Math. Jussieu*, 17(4):979–980, 2018. [1](#), [15](#)
- [CHS10] Steven Dale Cutkosky, Jürgen Herzog, and Hema Srinivasan. Asymptotic growth of algebras associated to powers of ideals. *Math. Proc. Cambridge Philos. Soc.*, 148(1):55–72, 2010. [9](#)
- [DW19] Omprokash Das and Joe Waldron. On the log minimal model program for 3-folds over imperfect fields of characteristic $p > 5$. *arXiv e-prints*, page arXiv:1911.04394, Nov 2019. [1](#), [16](#)
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52. [15](#)
- [Has10] Mitsuyasu Hashimoto. F -pure homomorphisms, strong F -regularity, and F -injectivity. *Comm. Algebra*, 38(12):4569–4596, 2010. [15](#)
- [HH89] Melvin Hochster and Craig Huneke. Tight closure and strong F -regularity. *Mém. Soc. Math. France (N.S.)*, (38):119–133, 1989. Colloque en l’honneur de Pierre Samuel (Orsay, 1987). [1](#)
- [HH90] Melvin Hochster and Craig Huneke. Tight closure, invariant theory, and the Briançon-Skoda theorem. *J. Amer. Math. Soc.*, 3(1):31–116, 1990. [1](#), [5](#), [12](#), [15](#)
- [HH91] Melvin Hochster and Craig Huneke. Tight closure and elements of small order in integral extensions. *J. Pure Appl. Algebra*, 71(2-3):233–247, 1991. [1](#)
- [HH93] Melvin Hochster and Craig Huneke. Phantom homology. *Mem. Amer. Math. Soc.*, 103(490):vi+91, 1993. [1](#)
- [HH94a] Melvin Hochster and Craig Huneke. F -regularity, test elements, and smooth base change. *Trans. Amer. Math. Soc.*, 346(1):1–62, 1994. [1](#), [15](#), [16](#)

- [HH94b] Melvin Hochster and Craig Huneke. Tight closure of parameter ideals and splitting in module-finite extensions. *J. Algebraic Geom.*, 3(4):599–670, 1994. [1](#), [15](#)
- [Hoa93] Le Tuan Hoa. Reduction numbers and Rees algebras of powers of an ideal. *Proc. Amer. Math. Soc.*, 119(2):415–422, 1993. [9](#), [15](#)
- [HS06] Craig Huneke and Irena Swanson. *Integral closure of ideals, rings, and modules*, volume 336 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2006. [6](#)
- [HS15] Craig Huneke and Ilya Smirnov. Prime filtrations of the powers of an ideal. *Bull. Lond. Math. Soc.*, 47(4):585–592, 2015. [9](#)
- [HW02] Nobuo Hara and Kei-Ichi Watanabe. F-regular and F-pure rings vs. log terminal and log canonical singularities. *J. Algebraic Geom.*, 11(2):363–392, 2002. [16](#)
- [ILL⁺07] Srikanth B. Iyengar, Graham J. Leuschke, Anton Leykin, Claudia Miller, Ezra Miller, Anurag K. Singh, and Uli Walther. *Twenty-four hours of local cohomology*, volume 87 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2007. [15](#)
- [KR86] Daniel Katz and Louis J. Ratliff, Jr. On the symbolic Rees ring of a primary ideal. *Comm. Algebra*, 14(5):959–970, 1986. [9](#)
- [Lip69] Joseph Lipman. Rational singularities, with applications to algebraic surfaces and unique factorization. *Inst. Hautes Études Sci. Publ. Math.*, (36):195–279, 1969. [1](#)
- [LS99] Gennady Lyubeznik and Karen E. Smith. Strong and weak F -regularity are equivalent for graded rings. *Amer. J. Math.*, 121(6):1279–1290, 1999. [1](#)
- [Mac96] Brian Cameron MacCrimmon. *Strong F -regularity and boundedness questions in tight closure*. ProQuest LLC, Ann Arbor, MI, 1996. Thesis (Ph.D.)—University of Michigan. [12](#)
- [MPST19] Linquan Ma, Thomas Polstra, Karl Schwede, and Kevin Tucker. F -signature under birational morphisms. *Forum Math. Sigma*, 7:e11, 20, 2019. [15](#)
- [PT18] Thomas Polstra and Kevin Tucker. F -signature and Hilbert-Kunz multiplicity: a combined approach and comparison. *Algebra Number Theory*, 12(1):61–97, 2018. [12](#)
- [Smi93] Karen Ellen Smith. *Tight closure of parameter ideals and F -rationality*. ProQuest LLC, Ann Arbor, MI, 1993. Thesis (Ph.D.)—University of Michigan. [12](#)
- [Smi97] Karen E. Smith. F -rational rings have rational singularities. *Amer. J. Math.*, 119(1):159–180, 1997. [1](#)
- [SS10] Karl Schwede and Karen E. Smith. Globally F -regular and log Fano varieties. *Adv. Math.*, 224(3):863–894, 2010. [16](#)
- [Tru98] Ngô Việt Trung. The Castelnuovo regularity of the Rees algebra and the associated graded ring. *Trans. Amer. Math. Soc.*, 350(7):2813–2832, 1998. [15](#)
- [Wat94] Keiichi Watanabe. Infinite cyclic covers of strongly F -regular rings. In *Commutative algebra: syzygies, multiplicities, and birational algebra (South Hadley, MA, 1992)*, volume 159 of *Contemp. Math.*, pages 423–432. Amer. Math. Soc., Providence, RI, 1994. [15](#)
- [Wil95] Lori J. Williams. Uniform stability of kernels of Koszul cohomology indexed by the Frobenius endomorphism. *J. Algebra*, 172(3):721–743, 1995. [1](#), [12](#), [16](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA
Email address: aberbachi@missouri.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22903 USA
Email address: tp2tt@virginia.edu